

The OCE package

Dan E. Kelley

March 31, 2008

Abstract

The `oce` package makes it easy to read, summarize and plot data from a variety of Oceanographic instruments, isolating the researcher from the quirky data formats that are common in this field. It also provides functions for working with basic seawater properties such as the equation of state, and with derived quantities such as the buoyancy frequency. Although simple enough to be used in a teaching context, `oce` is powerful enough for a research setting. These things are illustrated here with practical examples.

1 Introduction

Oceanographers must deal with measurements made by a wide variety of instruments, a task that is complicated by the delight instrument manufacturers seem to take in inventing new data formats. The manufacturers often provide software for scanning the data files and producing some standard plots, but this is of limited use to researchers who work with several instrument types at the same time, and who need to carry the analysis beyond the first step.

The need to scan diverse data files was one motivation for the creation of `oce`, but an equal goal was to make it easy to work with the data once they are in the system. This was accomplished partly by the provision of functions to work with the data, and partly by developing a uniform object design that lets users reach inside without guesswork¹, and partly by adhering to a policy of adding features according to the priorities of practical research. The net result is that `oce` is a fairly comfortable tool today, and one that should to remain so as it grows.

2 Calculations of seawater properties

The `oce` package provides many functions for dealing with seawater properties. Probably the most used is `sw.rho(S,t,p)`, which computes seawater density ρ as a function of salinity S (PSU), in-situ temperature t ($^{\circ}\text{C}$... note the use of lower-case in this and related functions, to avoid confusion with `T`, an abbreviation used sometimes in R programs) and pressure p (decibar). The result is a number in the order of 1000kg/m^3 . For many purposes, Oceanographers prefer to use the density anomaly $\sigma = \rho - 1000\text{kg/m}^3$, provided with `sw.sigma(S,t,p)`, or its adiabatic cousin σ_{θ} , provided with `sw.sigma.theta(S,t,p)`.

Note that the names of functions relating to seawater material properties start with “`sw.`”; future versions of `oce` may add similarly named functions for the properties of air, sediment, organisms, etc.

The `sw.` functions can also take a `ctd` object instead of the `(S, t, p)` arguments.

¹Each `oce` object is a list containing three elements: (a) `data`, a list or a data frame containing the actual data, for a CTD object, this will contain pressure, temperature, etc., (b) `metadata`, a list containing data such things as file headers, the location of a CTD cast, etc., and (c) `processing.log`, a list that documents how the file was created (often by a `read` or `as` method) and how it was changed thereafter (e.g. by decimating a CTD cast).

Most of the functions use the UNESCO formulations of seawater properties, but new formulations may be added as they come into use in the literature. A partial list of seawater functions is as follows: `sw.dynamic.height` (dynamic height), `sw.N2` (buoyancy frequency), `sw.S.C.T.p` (salinity S from C , T and p), `sw.S.T.rho` (S from T and ρ), `sw.T.S.rho` (T from S and ρ), `sw.T.freeze` (freezing temperature), `sw.alpha` (thermal expansion coefficient $\alpha = -\rho_0^{-1}\partial\rho/\partial T$), `sw.beta` (haline compression coefficient $\beta = \rho_0^{-1}\partial\rho/\partial S$), `sw.alpha.over.beta` (α/β), `sw.conductivity` (conductivity from S , T and p), `sw.depth` (depth from p and latitude), `sw.lapse.rate` (adiabatic lapse rate), `sw.rho` (density ρ from S , T and p), `sw.sigma` ($\rho - 1000 \text{ kg/m}^3$), `sw.sigma.t` (σ with p set to zero and temperature unaltered), `sw.sigma.theta` (σ with p set to zero and temperature altered adiabatically), `sw.sound.speed` (speed of sound in m/s), `sw.specific.heat` (specific heat in J/kg/°C), `sw.spice` (a quantity used in double-diffusive research), `sw.theta` (potential temperature in °C), and `sw.viscosity` (viscosity). Details and examples are, of course, provided in the documentation of these functions.

Exercise 1. (a) What is the density of a seawater parcel at pressure 100dbar, with salinity 34PSU and temperature 10°C? (b) What temperature would the parcel have if raised adiabatically to the surface? (c) What density would it have if raised adiabatically to the surface? (d) What density would it have if lowered about 100m, increasing the pressure to 200dbar? (e) Draw a blank TS diagram with S from 30 to 40PSU and T from -2 to 20°C . (Answers are provided at the end of this document.)

3 CTD data

3.1 Example with pre-trimmed data

To get you started with CTD data, `oce` provides a sample data set that has been trimmed to just the downcast portion of the sampling. (See the next section to learn how to do this trimming.). The commands

```
> library(oce)
> data(ctd)
> plot(ctd)
```

produce Figure 1. You may also get a summary of the data with

```
> summary(ctd)
```

The object used to hold CTD data stores not just the data, but also the raw header sequence, and whatever has been discovered about the dataset by parsing the header; use

```
> names(ctd)
```

to learn about these metadata, and use

```
> names(ctd$data)
```

to find out what sensors were attached to the instrument, thus providing data columns.

As noted above, the seawater-property functions (whose names begin with `sw.`) can take a `ctd` object instead of the individual arguments S , t , and p .

It is possible to plot the components of a `ctd` object individually, either by accessing the data directly (see the WOCE section, below, for an example) or by using more specialized functions such as `plot.TS` and `plot.profile`.

Exercise 2. Plot a profile of σ_θ and N^2 , for just the data in the pycnocline.




Figure 1: Overview graph of the sample CTD dataset `ctd`, acquired during the St Lawrence Estuary Internal Wave Experiment. (This dataset has been trimmed to just the downcast; see the text and Figure 2 for more on trimming.)

3.2 Example with raw data

Practicing Oceanographers may be wondering how the CTD cast used in the preceding section was trimmed of equilibration-phase and upcast-phase data. Data from these sections are spurious and must be trimmed as a first step in processing. For example, consider the following code.

```
> data(ctd.raw)
> plot.ctd.scan(ctd.raw)
```

This produces a two-panel plot (Figure 2) of the data as a time-series, revealing not just the (useful) downcast, but also the subsequent upcast sequence. The x-axis in this plot is the scan number, which is a convenient index for extraction of the downcast portion of the profile by an essentially manual method, e.g. proceeding with a sequence of commands such as

```
> plot.ctd.scan(ctd.trim(ctd.raw, "scan", c(140, 250)))
> plot.ctd.scan(ctd.trim(ctd.raw, "scan", c(150, 250)))
```

This is the “gold standard” method, which is recommended for detailed work. However, for quick work, you may find that the automatic downcast detection scheme works adequately, e.g.

```
> ctd.trimmed <- ctd.trim(ctd.raw)
```

It should be noted that `ctd.trim` inserts entries into the object’s log file, so that you (or anyone else with whom you share the object) will be able to see the details of how the trimming was done.

Once the profile has been trimmed, you may wish to use `ctd.decimate()` to smooth the data and interpolate the smoothed results to uniformly-spaced pressure values. For example, a quick examination of a file might be done by the following:

```
> plot(ctd.decimate(ctd.trim(read.ctd("stn123.cnv"))))
```




Figure 2: Scanwise plot of the `ctd.raw` sample data set. Note the wild spike at the start, the equilibration phase before the downcast, and the spurious freshening signal near the start of the upcast. See the text for a discussion of how inspection of such graphs can help in trimming CTD data.

3.3 Example with WOCE archive data

The package has a harder time scanning the headers of data files in the WOCE archive format than it does in the Seabird format illustrated in the previous examples. This is mainly because front-line researchers tend to work in the Seabird format, and partly because the WOCE format is odd. For example, the first line of a WOCE file is of the form `CTD,20060609WHPOSIODAM`. Only the first part of this is easy to scan. To the left of the comma is the string `CTD`, which seems straightforward, except that in some cases it is written `CTD0`. After the comma is found the file date (`yyyymmdd`), and that is easy to parse. But then things get tricky. The next sequence is a string of characters that indicate the division of the institute (`WHPO`), the institute itself (`SIO`), and the name of the investigator (`DAM`). The problem is that no dividers separate these items, and that there are no standards for the item lengths. The approach to `oce` development is to tackle easy and important problems before complicated special cases, and so no attempt has been made to parse this part of the header. Of course, R provides access to object constituents, so that a human working with this file could easily do e.g.

```
> x <- read.ctd("nnsa_00934_00001_ct1.csv", type = "WOCE")
> x$metadata$institute <- "SIO"
> x$metadata$scientist <- "DAM"
```

For a real-world example (with warts!), visit http://cchdo.ucsd.edu/data_access?ExpoCode=58JH199410 and download the zip file containing the Arctic section called “CARINA”, measured in 1994. Expand the zip file, enter the directory, and run the code below.

```
> library(oce)
> files <- system("ls *.csv", intern = TRUE)
> for (i in 1:length(files)) {
+   cat(files[i], "\n")
+   x <- read.ctd(files[i])
+   if (i == 1) {
```

```

+         plot.TS(x, xlim = c(31, 35.5), ylim = c(-1, 10), type = "l",
+               col = "red")
+     }
+     else {
+         lines(x$data$salinity, x$data$temperature, col = "red")
+     }
+ }

```

What you'll see is an overall $T-S$ diagram for the entire dataset. It may take a while, since the dataset contains over 90,000 observations. You may note that, even though this is an official, quality-controlled dataset, it is not without problems. The graph that is produced by this code has several spurious lines oriented horizontally (indicating spurious salinity) and vertically (indicating spurious temperature). One way to find such values is to put the lines

```

> print(range(x$data$temperature))
> print(range(x$data$salinity))

```

after the `read.ctd()` command. One thing you'll find is that station 987 has a minimum salinity range of 0.0009 to 987. These values are clearly in error, as are the temperatures at this spot in the file. (It is perhaps revealing that the spurious salinity is equal to the station number.) Indeed, at this spot in the file it can be seen that the pressure jumps from 1342 to 0, and then starts increasing again; the file contains two profiles, or the same profile twice. This is not the only flaw that is revealed by the graph, and by `range` commands; a generous user would spend a week tracking down such issues, and would then contact the data provider (or the chief scientist of the field work) with specific suggestions for correcting the files. The point here is to highlight how this package can be used with real-world data.

3.4 Section plots

The commands

```

> data(section)
> data(coastline.hal)
> plot(section, coastline = coastline.hal)

```

will plot a summary diagram (Figure 3) containing sections of T , S , and σ_θ , along with a chart indicating station locations. In addition to such overview diagrams, `plot` can also create individual plots of individual properties.

Exercise 3. Draw a TS diagram for the section data, colour-coded by station

The Halifax section is supplied in a pre-gridded format, but some datasets have different pressure levels at each station. For such cases, the `section.grid` function may be used, e.g.

```

> data(a03)
> Gulf.Stream <- section.subset(a03, 124:102)
> Gulf.Stream.gridded <- section.grid(Gulf.Stream, p = seq(0, 5000,
+   10))
> data(coastline.world)
> plot(Gulf.Stream.gridded, coastline = coastline.world, map.xlim = c(-80,
+   -60))

```




Figure 3: CTD section of data acquired in Halifax Harbour in the autumn of 2003 by students in the *Introduction to Physical Oceanography* class at Dalhousie University. (Dan Kelley teaches this class, and the work at sea was supervised by his teaching assistant Natacha Bernier, who received her PhD from Dalhousie in 2005.) The stations go from a site near the Sackville river inflow at the northwest portion of Bedford Basin to site at the entrance of the harbour. (The author's office at Dalhousie is located near the bottom axis at 63.58W.)




Figure 4: Portion of the CTD section designated A03 (Cruise chief scientist: Tereschenkov, SOI.), showing the region of the Gulf Stream. (The small number of S contours in the upper water column is the result of a near-bottom salinity anomaly in the station at about 200km.)




Figure 5: Coastline of eastern Canada, showing Prince Edward Island, New Brunswick, and Nova Scotia. The blue circle indicates the location of Halifax, the capital of Nova Scotia.

produces Figure 4. The ship doing the sampling was travelling westward from the Mediterranean towards North America, taking 124 stations in total; the `station.indices` value selects the last few stations of the section, during which the ship heading was changed to run in a northwesterly direction, to cross isobaths (and perhaps, the Gulf Stream) at right angles.

Exercise 4. Plot dynamic height across the Gulf Stream, and show the corresponding geostrophic velocity.

4 Coastline data

The commands

```
> library(oce)
> data(coastline.maritimes)
> plot(coastline.maritimes, col = "darkred")
> points(-(63 + 34/60), 44 + 39/60, cex = 3, col = "blue")
```

produce a map of the coastline of Eastern Canada (Figure 5). Such coastline data are available from a variety of sources. The NOAA site http://www.ngdc.noaa.gov/mgg_coastline/ is particularly popular, and it has the advantage of providing data in Splus format. The function `read.coastline` can handle reading that format (plus some other formats).




Figure 6: Sea-level timeseries measured in 2003 in Halifax Harbour. (The spike in September is the storm surge associated with Hurricane Juan, regarded by the Canadian Hurricane Centre to be one of the most powerful and damaging hurricanes to ever hit Canada.)

5 Sea-level data

5.1 Time-domain analysis

The commands

```
> library(oce)
> data(sealevel.hal)
> plot(sealevel.hal)
```

load and graph a build-in dataset of sea-level timeseries. The result, shown in Figure 6, is a four-panel plot. The top panel is a timeseries view that provides an overview of the entire data set. The second panel is narrowed to the most recent month, which should reveal spring-neap cycles if the tide is mixed. The third panel is a spectrum, with a few tidal constituents indicated. At the bottom is a cumulative spectrum, which makes these narrow-banded constituents quite visible.

Exercise 5. Illustrate Halifax sealevel variations during Hurricane Juan

Exercise 6. Draw a spectrum of sea-level variation, with the M2 tidal component indicated.




Figure 7: Sea-level variation in Halifax Harbour in 2003 (black line) and the residual after removal of tides using `tidem()`. The spike on September 28th is a storm surge caused by Hurricane Juan, which caused a great deal of damage to the region.

5.2 Tidal analysis

Tidal analysis is provided in a rudimentary way, as is illustrated in the example below, which shows the storm surge experienced in Halifax in 2003, with Hurricane Juan.

```
> library(oce)
> data(sealevel.hal)
> tide <- tidem(sealevel.hal)
> days <- 15
> n <- length(sealevel.hal$data$eta)
> i <- seq(n - 24 * days, n)
> t <- sealevel.hal$data$t[i]
> eta <- sealevel.hal$data$eta[i]
> eta.pred <- predict(tide)[i]
> plot(t, eta, type = "l", ylim = c(-0.5, 3))
> abline(h = 0, col = "pink")
> lines(t, eta - eta.pred, col = "red")
```

produce Figure 7, illustrating the use of tidal analysis to detect storm surges. Note the use of the `model` that is returned by the tide analysis.

(It should be noted that the tidal analysis portion of `oce` is still in active development, and the features are subject to change. Use `?tidem` to learn more about this.)




Figure 8: Lobo measurements in Northwest Arm of Halifax Harbour, at the time of the 2007 Spring bloom.

6 Lobo data

The commands

```
> library(oce)
> data(lobo)
> plot(lobo)
```

produce a plot (Figure 8) of lobo data from the Northwest Arm of Halifax Harbour. Note the relationship between decreasing nutrients and increasing fluorescence, as well as the diurnal signal in the latter.

The reader should note that the `lobo` part of `oce` is somewhat preliminary. In particular, the package requires that certain data columns be present, and in a certain order. Also, the function `read.oce` does not understand `lobo` files. Why these limitations, you ask? Well, the `lobo` code was really only written as an aside, for the author’s contribution to a “predict the spring bloom” contest held at Dalhousie University.

Exercise 7. Draw a $T - S$ plot for these data, using a colour coding to indicate time, and using plotting tricks to reduce the obscuring of this time signal.

7 The future of oce

The present version of `oce` can only handle data types that the author has been using lately in his research. New data types will be added as the need arises in that work, but the author would also be happy to add other data types that are likely to prove useful to the Oceanographic community. (The data types need not be restricted to Physical Oceanography, but the author will need some help in dealing with other types of data, given his research focus.)

As for algorithms, there are plenty of gaps in `oce`. Dealing with measurements of turbulence is a high priority for the author, for example, and it is also clear that there are some methods of ADP processing that should be provided by the package.

Two principles will guide the addition of data types and functions: (a) the need, as perceived by the author or by other contributors and (b) the ease with which the additions can be made.

The site <http://code.google.com/p/r-oce/> provides a window on the development that goes on between the CRAN releases of the package. Please visit the site to report bugs, to suggest new features, or just to see how `oce` development is coming along.

Answers to exercises

Exercise 1 – Seawater properties.

```
> library(oce)
> sw.rho(S = 34, t = 10, p = 100)

[1] 1026.624

> sw.theta(S = 34, t = 10, p = 100)

[1] 9.988598

> sw.rho(S = 34, t = sw.theta(S = 34, t = 10, p = 100), p = 0)

[1] 1026.173

> sw.rho(S = 34, t = sw.theta(S = 34, t = 10, p = 100, pref = 200),
+       p = 200)

[1] 1027.074

> plot.TS(as.ctd(c(30, 40), c(-2, 20), rep(0, 2)), grid = TRUE,
+       col = "white")
```

Exercise 2 – Profile plots. Although one may argue as to the limits of the pycnocline, for illustration let us say it is in 5bar to 12dbar range.

```
> library(oce)
> data(ctd)
> pycnocline <- ctd.trim(ctd, "pressure", c(5, 12))
> plot.profile(pycnocline, type = "density+N2")
```

Exercise 3 – TS diagram for section data.

```
> library(oce)
> data(section)
> SS <- TT <- pp <- id <- NULL
> n <- length(section$data$station)
> for (i in 1:n) {
+   stn <- section$data$station[[i]]
+   SS <- c(SS, stn$data$salinity)
+   TT <- c(TT, stn$data$temperature)
+   pp <- c(pp, stn$data$pressure)
+   id <- c(id, rep(i, length(stn$data$pressure)))
+ }
> ctd <- as.ctd(SS, TT, pp)
> plot.TS(ctd, col = hsv(0.7 * id/n), cex = 2, pch = 21)
```

Exercise 4 – Gulf Stream. (Try ?sw.dynamic.height for hints on smoothing.)

```
> library(oce)
> data(a03)
> Gulf.Stream <- section.subset(a03, 124:102)
> dh <- sw.dynamic.height(Gulf.Stream)
> par(mfrow = c(2, 1))
> plot(dh$distance, dh$height, type = "b", xlab = "", ylab = "Dyn. Height [m]")
> grid()
> f <- coriolis(Gulf.Stream$data$station[[1]]$metadata$latitude)
> g <- gravity(Gulf.Stream$data$station[[1]]$metadata$latitude)
> v <- diff(dh$height)/diff(dh$distance) * g/f/1000
> plot(dh$distance[-1], v, type = "l", col = "blue", xlab = "Distance [km]",
+   ylab = "Velocity [m/s]")
> grid()
> abline(h = 0)
```

Exercise 5 – Halifax sealevel during Hurricane Juan. A web search will tell you that Hurricane Juan hit about midnight, 2003-sep-28. The author can verify that the strongest winds occurred a bit after midnight – that was the time he moved to a room without windows, in fear of flying glass.

```
> library(oce)
> data(sealevel.hal)
> plot(sealevel.hal, focus.time = c("2003-09-23", "2003-10-05"))
> abline(v = as.POSIXct("2003-09-28 23:30:00"), col = "red", lty = "dotted")
> mtext("Hurricane\nJuan", at = as.POSIXct("2003-09-28 23:30:00"),
+   col = "red")
```

Exercise 6 – Sealevel spectrum. Notice the use of (object)\$data\$(item) here. All oce objects are lists, and all of them contain a data element of a similar form to this.

```
> library(oce)
> data(sealevel.hal)
> spectrum(sealevel.hal$data$eta, spans = c(3, 7))
> abline(v = 1/12.42)
> mtext("M2", at = 1/12.42, side = 3)
```

Exercise 7 – Lobo plot. The resampling with `i` is to avoid obscuring colours by overplotting. Note the use of `as.ctd` to assemble the data into something that `plot.TS` can handle. This is an example of the practicality of `oce`; eventually, `plot.TS` may be altered to take simple columns of data, but for now it seems reasonable to require the user to assemble these data into a CTD object, and to spend development time on something that will pay off better.

```
> library(oce)
> data(lobo)
> i <- sample(length(lobo$data$temperature))
> a <- as.numeric(lobo$data$time[i] - lobo$data$time[1])
> col <- hsv(0.5 * a/max(a), 1, 1)
> plot.TS(as.ctd(lobo$data$salinity[i], lobo$data$temperature[i],
+               0), col = col, pch = 1)
```

Index

calculation

- seawater properties, 1

data

- coastline, 8

- lobo timeseries, 14

- dynamic height, 1

Gulf Stream, 5

- geostrophic calculation, 13

Hurricane Juan

- surge detection by detiding, 10

- surge seen in time-series of sea level, 9

- worked example of sea-level plot, 13

reading

- ctd profile, 3

- ctd section, 5

sea level

- during Hurricane Juan, 9, 13

- storm surge during Hurricane Juan, 10

- seawater properties, calculations of, 1

section

- extracting profile data from, 12