

ARMA-GARCH modelling and white noise tests

James Proberts
University of Manchester

Georgi N. Boshnakov
University of Manchester

Abstract

This vignette illustrates applications of white noise tests in GARCH modelling. It is based on an example from an MMath project by the first author.

Keywords: autocorrelations, white noise tests, IID tests, GARCH models, time series.

1. The data

In this example we consider data from Freddie Mac, a mortgage loan company in the USA. This stock is an interesting case for study. In the financial crash of 2008 it dropped from roughly \$60 to \$0.5 over the course of a year. It is now (April 2017) majority owned by the government and has all its profits and dividends swepted. There has been speculation on this stock being returned to private ownership for years making it prone to clusters of volatility. We import weekly data from Yahoo Finance covering the period from 10/05/2006 to 22/04/2017, and calculate the weekly simple log returns.

```
R> ## using a saved object, originally imported with:  
R> ## FMCC <- yahooSeries("FMCC", from = "2006-05-10", to = "2017-04-22",  
R> ##                               freq = "weekly")  
R> FMCC <- readRDS("FMCC.rds")  
R> logreturns <- diff(rev(log(FMCC$FMCC.Close)))
```

A plot of the log-returns. is given in Fig. 1. We also calculate the autocorrelations and partial autocorrelations for the log returns.

```
R> FMCClr.acf <- autocorrelations(logreturns)  
R> FMCClr.pacf <- partialAutocorrelations(logreturns)
```

2. Autocorrelations

We now produce a plot of the autocorrelations to assess whether the series is autocorrelated, see Fig. 2. There are two bounds plotted on the graph. The straight red line represents the standard bounds under the strong white noise assumption. The second line is under the hypothesis that the process is GARCH.

Several autocorrelations seem significant under the iid hypothesis. This may lead us to fitting an ARMA or ARMA-GARCH model. On the other hand, the autocorrelations are well into

```
R> plot(logreturns, type="l", main="Log-returns of FMCC")
```




Figure 1: Log-returns of weekly log-returns of FMCC from 10 May 2006 to 22 Apr 2017.

the bands produced under the GARCH hypothesis, suggesting a pure GARCH model, without any ARMA terms. So, it matters on which test we base our decision.

The partial autocorrelation function can be used instead of the autocorrelations, with similar inferences, see Fig. 3.

3. Pormanteau tests

Routine portmanteau tests, such as Ljung-Box, also reject the IID hypothesis. Here we carry out IID tests using the method of Li-McLeod:

```
R> wntLM <- whiteNoiseTest(FMCClr.acf, h0 = "iid", nlags = c(5,10,20),
+                           x = logreturns, method = "LiMcLeod")
R> wntLM$test
```

	ChiSq	DF	pvalue
[1,]	37.18469	5	5.499929e-07

```
R> plot(FMCClr.acf, data = logreturns)
```




Figure 2: Autocorrelation test of the log returns of FMCC

```
[2,] 76.99131 10 1.946524e-12
[3,] 103.19392 20 3.363466e-13
attr("method")
[1] "LiMcLeod"
```


Small p-values lead to rejection of the null hypothesis at reasonable levels. Rejection of the null hypothesis is often taken to mean that the data are autocorrelated.

Let us test for fitting a GARCH-type model by using the following code which has the weaker assumption that the log returns are GARCH. Let us change the null hypothesis to "garch" (one possible weak white noise hypothesis):

```
R> wntg <- whiteNoiseTest(FMCClr.acf, h0 = "garch", nlags = c(5,10,15), x = logreturns)
R> wntg$test
```

```
      h      Q      pval
[1,]  5  5.115077 0.4019983
```

```
R> plot(FMCClr.pacf, data = logreturns,  
+ main="Partial Autocorrelation test of the log returns of FMCC")
```



```
[2,] 10 12.667143 0.2428821
[3,] 15 18.036831 0.2607342
```

The high p-values give no reason to reject the hypothesis that the log-returns are a GARCH white noise process. In other words, there is no need to ARMA modelling.

4. Fitting GARCH(1,1) models and their variants

Based on the discussion above, we go on to fit GARCH model(s), starting with a GARCH(1,1) model with Gaussian innovations.

```
R> fit1 <- garchFit(~garch(1,1), data = logreturns, trace = FALSE)
R> summary(fit1)
```

```
Title:
  GARCH Modelling
```

```
Call:
  garchFit(formula = ~garch(1, 1), data = logreturns, trace = FALSE)
```

```
Mean and Variance Equation:
  data ~ garch(1, 1)
<environment: 0x000000000c3cf670>
  [data = logreturns]
```

```
Conditional Distribution:
  norm
```

```
Coefficient(s):
      mu      omega    alpha1    beta1
-6.3541e-05  2.9206e-03  4.3649e-01  5.8992e-01
```

```
Std. Errors:
  based on Hessian
```

```
Error Analysis:
      Estimate Std. Error t value Pr(>|t|)
mu      -6.354e-05  5.006e-03  -0.013    0.99
omega    2.921e-03  6.982e-04   4.183 2.87e-05 ***
alpha1   4.365e-01  7.623e-02   5.726 1.03e-08 ***
beta1    5.899e-01  5.427e-02  10.869 < 2e-16 ***
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Log Likelihood:
 341.7229    normalized: 0.5995139
```

Description:

Mon Jul 09 09:44:51 2018 by user: mcbssgb2

Standardised Residuals Tests:

			Statistic	p-Value
Jarque-Bera Test	R	Chi ²	24230.08	0
Shapiro-Wilk Test	R	W	0.7433933	0
Ljung-Box Test	R	Q(10)	9.525801	0.4830325
Ljung-Box Test	R	Q(15)	12.92386	0.6081792
Ljung-Box Test	R	Q(20)	14.75224	0.7904048
Ljung-Box Test	R ²	Q(10)	0.7315935	0.9999597
Ljung-Box Test	R ²	Q(15)	0.9445704	0.9999998
Ljung-Box Test	R ²	Q(20)	1.338934	1
LM Arch Test	R	TR ²	0.8791397	0.9999931

Information Criterion Statistics:

AIC	BIC	SIC	HQIC
-1.184993	-1.154497	-1.185090	-1.173094

The diagnostics suggest that the standardised residuals and their squares are IID and that the ARCH effects have been accommodated by the model. Their distribution is clearly not Gaussian however (see the p-values for Jarque-Bera and Shapiro-Wilk Tests), so another conditional distribution can be tried.

Another possible problem is that $\alpha_1 + \beta_1 > 0$.

```
R> fit2 <- garchFit(~garch(1,1), cond.dist = c("sstd"), data = logreturns, trace = FALSE)
R> summary(fit2)
```

Title:

GARCH Modelling

Call:

```
garchFit(formula = ~garch(1, 1), data = logreturns, cond.dist = c("sstd"),
  trace = FALSE)
```

Mean and Variance Equation:

```
data ~ garch(1, 1)
<environment: 0x0000000011fb12c8>
[data = logreturns]
```

Conditional Distribution:

sstd

Coefficient(s):

mu	omega	alpha1	beta1	skew	shape
0.00024523	0.00277227	0.99999999	0.73057510	1.16531856	2.14375224

Std. Errors:
based on Hessian

Error Analysis:

	Estimate	Std. Error	t value	Pr(> t)
mu	0.0002452	0.0033295	0.074	0.9413
omega	0.0027723	0.0017142	1.617	0.1058
alpha1	1.0000000	0.5302728	1.886	0.0593 .
beta1	0.7305751	0.0763615	9.567	<2e-16 ***
skew	1.1653186	0.0576821	20.202	<2e-16 ***
shape	2.1437522	0.0969271	22.117	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Log Likelihood:

555.2528 normalized: 0.9741278

Description:

Mon Jul 09 09:44:52 2018 by user: mcbssgb2

Standardised Residuals Tests:


			Statistic	p-Value
Jarque-Bera Test	R	Chi^2	27547.01	0
Shapiro-Wilk Test	R	W	0.7324028	0
Ljung-Box Test	R	Q(10)	7.820836	0.6463324
Ljung-Box Test	R	Q(15)	10.34984	0.7971759
Ljung-Box Test	R	Q(20)	11.87712	0.9202405
Ljung-Box Test	R^2	Q(10)	0.7097748	0.9999651
Ljung-Box Test	R^2	Q(15)	1.089078	0.9999995
Ljung-Box Test	R^2	Q(20)	1.449253	1
LM Arch Test	R	TR^2	0.9024198	0.999992

Information Criterion Statistics:

AIC	BIC	SIC	HQIC
-1.927203	-1.881459	-1.927421	-1.909355

The qq-plot of the standardised residuals, suggests that the fitted standardised skew-t conditional distribution is not good enough.

R> plot(fit2, which = 13)



```
R> fit3 <- garchFit(~aparch(1,1), cond.dist = c("sstd"), data = logreturns, trace = FALSE)
R> summary(fit3)
```

Title:

GARCH Modelling

Call:

```
garchFit(formula = ~aparch(1, 1), data = logreturns, cond.dist = c("sstd"),
  trace = FALSE)
```

Mean and Variance Equation:

```
data ~ aparch(1, 1)
```

```
<environment: 0x00000000e42c520>
```

```
[data = logreturns]
```

Conditional Distribution:

```
sstd
```

Coefficient(s):

mu	omega	alpha1	gamma1	beta1	delta	skew
----	-------	--------	--------	-------	-------	------


```
0.0034733  0.0425723  1.0000000  0.2025020  0.7970743  0.7374463  1.2050209
      shape
2.0099078
```

```
Std. Errors:
  based on Hessian
```

```
Error Analysis:
```

	Estimate	Std. Error	t value	Pr(> t)
mu	0.003473	0.002574	1.349	0.177211
omega	0.042572	0.020910	2.036	0.041752 *
alpha1	1.000000	0.520738	1.920	0.054814 .
gamma1	0.202502	0.147550	1.372	0.169930
beta1	0.797074	0.045315	17.590	< 2e-16 ***
delta	0.737446	0.210335	3.506	0.000455 ***
skew	1.205021	0.052296	23.042	< 2e-16 ***
shape	2.009908	0.004466	450.084	< 2e-16 ***

```
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Log Likelihood:
```

```
557.6893      normalized:  0.9784022
```

```
Description:
```

```
Mon Jul 09 09:44:53 2018 by user: mcbssgb2
```

```
Standardised Residuals Tests:
```

			Statistic	p-Value
Jarque-Bera Test	R	Chi^2	17899.48	0
Shapiro-Wilk Test	R	W	0.7489928	0
Ljung-Box Test	R	Q(10)	10.27713	0.4165255
Ljung-Box Test	R	Q(15)	13.3245	0.5772494
Ljung-Box Test	R	Q(20)	15.23641	0.7627242
Ljung-Box Test	R^2	Q(10)	2.251332	0.9940257
Ljung-Box Test	R^2	Q(15)	2.625092	0.9998262
Ljung-Box Test	R^2	Q(20)	3.271614	0.9999914
LM Arch Test	R	TR^2	2.218819	0.9989894

```
Information Criterion Statistics:
```

AIC	BIC	SIC	HQIC
-1.928734	-1.867743	-1.929121	-1.904937

The qq-plots of the standardised results for all models fitted above suggest that the chosen conditional distributions are unsatisfactory. Moreover, the fitted standardised-t distributions have shape parameters (degrees of freedom) slightly over two. Suggesting extremely heavy tails, maybe even the need for stable distributions.

Note also that in all models above $\alpha_1 + \beta_1$ is greater than one, a possible violation of any form of stationarity.

Or maybe, it is just that the GARCH models tried here are not able to accomodate varying behaviour before, during and after the financial crisis.

Affiliation:

Georgi N. Boshnakov

School of Mathematics

The University of Manchester

Oxford Road, Manchester M13 9PL, UK

URL: <http://www.maths.manchester.ac.uk/~gb/>